

Review

Open Access



Autophagy in ageing and ageing-related neurodegenerative diseases

Cansu Karabiyik^{1,2}, Rebecca A. Frake³, So Jung Park^{1,2}, Mariana Pavel⁴, David C. Rubinsztein^{1,2}

¹Department of Medical Genetics, Cambridge Institute for Medical Research, University of Cambridge, Cambridge CB2 0XY, UK.

²UK Dementia Research Institute, University of Cambridge, Cambridge CB2 0XY, UK.

³Centre for Discovery Brain Sciences, The University of Edinburgh, Edinburgh EH8 9XD, UK.

⁴Department of Immunology, Grigore T. Popa University of Medicine and Pharmacy, Iasi 700115, Romania.

Correspondence to: Prof. David C. Rubinsztein, Department of Medical Genetics, Cambridge Institute for Medical Research, University of Cambridge, The Keith Peters Building, Hills Road, Cambridge CB2 0XY, UK. E-mail: dcr1000@cam.ac.uk

How to cite this article: Karabiyik C, Frake RA, Park SJ, Pavel M, Rubinsztein DC. Autophagy in ageing and ageing-related neurodegenerative diseases. *Ageing Neur Dis* 2021;1:2. <https://dx.doi.org/10.20517/and.2021.05>

Received: 9 Jun 2021 **First Decision:** 30 Jun 2021 **Revised:** 6 Jul 2021 **Accepted:** 12 Jul 2021 **First online:** 14 Jul 2021

Academic Editor: Weidong Le **Copy Editor:** Yue-Yue Zhang **Production Editor:** Yue-Yue Zhang

Abstract

Autophagy is a catabolic mechanism that allows cells to deliver cytoplasmic contents to lysosomes for degradation to maintain energy homeostasis and to protect cells against stress. Autophagy has been directly linked to neurodegeneration and ageing by an extensive body of research. It has become evident that disruption of autophagy contributes significantly to age-related pathologies and to the cognitive and motor declines associated with “healthy” ageing. Autophagic dysfunction causes the accumulation of many of the toxic, aggregate-prone proteins that are responsible for neurodegenerative diseases, including mutant huntingtin, alpha-synuclein, tau, and others. Since upregulation of autophagy has been found to reduce levels of such protein species, the therapeutic potential of autophagy induction as a strategy against age-related diseases and a method for modulating longevity has been widely studied. Here we review the evidence supporting a role for autophagy dysfunction in the progression of the age-associated functional decline in the brain and age-related brain pathologies and discuss the available evidence that upregulation of autophagy may be a valuable therapeutic strategy.

Keywords: Autophagy, ageing, neurodegenerative diseases, misfolded proteins, protein degradation



© The Author(s) 2021. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, sharing, adaptation, distribution and reproduction in any medium or format, for any purpose, even commercially, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.



INTRODUCTION

Autophagy is a term used to describe three pathways that deliver cytoplasmic contents to lysosomes for digestion and recycling. These include microautophagy, chaperone-mediated autophagy (CMA), and macroautophagy. The three pathways coexist in most mammalian cells and differ in their modes of delivery of cytoplasmic cargo to the lysosomal compartment. In brief, microautophagy involves direct delivery of cytosolic components to the lysosome by invagination of the lysosomal membrane^[1], while CMA involves selective translocation of cytoplasmic proteins with a KFERQ peptide motif into the lysosomal lumen for degradation^[2]. In macroautophagy, the cytoplasmic cargo, which can include proteins and organelles, is sequestered and engulfed by double-membraned, cup-shaped structures known as phagophores that elongate and extend around the intracellular content to form an autophagosome. Autophagosomes ultimately fuse with lysosomes to become autolysosomes, after which the degradative enzymes degrade the autophagic contents. In this review, we will focus on macroautophagy (hereafter referred to as autophagy).

Autophagy can be divided into selective and non-selective forms of autophagy^[3]. Non-selective autophagy (often referred to as bulk autophagy) is considered in lacking cargo specificity and can be induced by various cellular stimuli, such as energy and nutrient deprivation. Selective autophagy increases the likelihood of cargo capture and can degrade aggregate-prone proteins (aggrephagy), intracellular pathogens (xenophagy), damaged mitochondria (mitophagy) and lysosomes (lysophagy), excess peroxisomes (pexophagy), and dysfunctional endoplasmic reticulum (ER-phagy).

Since it was first discovered in yeast as a survival mechanism^[4], it has become evident that autophagy is important for a variety of biological processes. Interestingly, perturbation of both selective and non-selective forms of autophagy has been associated with ageing and numerous age-related disorders, such as inflammation, cancer, diabetes and neurodegenerative diseases^[5].

The susceptibility of neurons to accumulate defective organelles and proteins due to their post-mitotic nature and the age-related decline in autophagic capacity appears to create an unfortunate paradox that puts autophagic dysfunction at the center of many neurodegenerative diseases where accumulation of toxic aggregated proteins is a hallmark and a causal factor for the disease pathogenesis'. Additionally, variants/polymorphisms in autophagy genes have been implicated in numerous age-related neurodegenerative diseases, including Alzheimer's disease (AD), Parkinson's disease (PD), Huntington's disease (HD) and amyotrophic lateral sclerosis/frontotemporal dementia (ALS/FTD) and may contribute to the accumulation of toxic intracellular protein aggregates in the brain. This has led to interesting questions regarding the potential of autophagy induction as a treatment against age-related diseases.

In this review, we will describe the evidence of autophagy dysfunction in ageing and age-related disorders and the potential of autophagy upregulation as a therapeutic strategy for the functional decline associated with ageing and neurodegenerative disorders.

AUTOPHAGY AND NEURODEGENERATIVE DISEASES

Intracellular protein misfolding and aggregation are common features of the abovementioned late-onset neurodegenerative diseases, also referred to as proteinopathies. The Mendelian mutations that cause proteinopathies frequently lead to toxic gain-of-functions in the disease-associated proteins, the expression of which correlate with disease severity. It has become increasingly evident that a given proteinopathy can be caused by various mutations that disturb different parts of the autophagic machinery [Figure 1 and Table 1]. Thus, it is critical to understand the cellular impact of the mutations that cause age-related neurodegenerative diseases. Additionally, since the toxic aggregated proteins associated with the

Table 1. Selected pathophysiological mechanisms related to autophagy impairment in AD, PD, HD and ALS

	Autophagosome formation	Cargo recruitment	AP-lysosome fusion	Lysosome function
AD	<ul style="list-style-type: none"> · PICALM depletion in AD brains inhibits autophagosome formation · Reduced Beclin-1 levels cause decreased autophagosome biogenesis · CCT complex impairment reduces autophagosome formation 	<ul style="list-style-type: none"> · PICALM depletion inhibits cargo recognition 	<ul style="list-style-type: none"> · Loss-of-function of PICALM impairs the fusion of autophagosomes with lysosomes in a VAMP8-dependent process · Reduced CCT/TRiC complex expression in AD brains blocks autophagosome delivery · Tau accumulation disturbs the integrity of lysosome membranes and impairs autophagosome maturation by inhibiting HDAC6 	<ul style="list-style-type: none"> · Decreased CCT/TRiC complex level in AD brains blocks lysosomal degradation · APOE4 variant increases Rab5-mediated endocytosis and causes non-degraded lipids and proteins to accumulate in swollen lysosomes.
PD	<ul style="list-style-type: none"> · VPS35 mutation causes ATG9 mislocalisation and inhibits autophagosome formation · Overexpression of α-syn <i>in vitro</i> and <i>in vivo</i> results in compromised autophagosome biogenesis through Rab1 inhibition, causing mislocalisation of ATG9 		<ul style="list-style-type: none"> · Inclusion bodies containing α-syn inhibit autophagosome maturation and fusion with lysosomes, resulting in decreased protein degradation 	<ul style="list-style-type: none"> · Depletion of ATP13A2 leads to decreased SYT11, which results in lysosomal dysfunction · iPSC-derived neurons from PD patients with GBA mutations showed increased α-syn levels, as well as autophagic and lysosomal defects · Accumulation of α-syn aggregates in the nervous system causes lysosomal impairment · PD mutation in LRRK2 G2019S induces defective lysosomal positioning mediated by Rab7 and enlarged lysosomes containing nondegraded contents
HD	<ul style="list-style-type: none"> · mHTT impairs autophagosome biogenesis 	<ul style="list-style-type: none"> · mHTT causes defects in cargo recognition by inhibiting organelle sequestration 	<ul style="list-style-type: none"> · mHTT perturbs post-Golgi trafficking to lysosomes, leading to dysfunctional autophagosome/lysosome dynamics 	
ALS		<ul style="list-style-type: none"> · ALS-associated L341V p62 exhibits defective binding to the core autophagy protein LC3 and impaired recruitment to autophagosomes · ALS/FTD-causing mutations of TBK1 impair binding and phosphorylation of optineurin, leading to reduced autophagy substrate recognition and clearance 	<ul style="list-style-type: none"> · Truncated CHMP2B blocks autophagy flux in cultured primary neurons from rat cortex and photoreceptor neurons in <i>Drosophila</i> eye sections · R155H mutation of VCP disrupts clearance of damaged lysosomes by autophagy 	
Ageing	<ul style="list-style-type: none"> · Reduced autophagosome formation in liver tissue from aged mice 		<ul style="list-style-type: none"> · Aged <i>C. elegans</i> has a dysfunctional late autophagic flux 	




Figure 1. Disruption of autophagy machinery in neurodegenerative diseases. Numerous genes implicated in neurodegenerative diseases such as AD, HD, PD and ALS have been associated with autophagy. These genes disrupt different parts of the autophagic machinery, including autophagosome formation, cargo recruitment, lysosomal function or autolysosomal fusion.

proteinopathies are substrates for autophagy, upregulation of autophagy or prevention of the age-related functional decline in autophagy may prove to be valuable as a therapeutic strategy.

Alzheimer's disease

AD is a progressive neurodegenerative disorder characterized by two main pathological hallmarks: the extracellular deposit of amyloid- β ($A\beta$) plaques and the intracellular accumulation of hyperphosphorylated tau tangles^[6]. There is a complex interplay between $A\beta$ and autophagy, as autophagy controls both the clearance and generation of $A\beta$. On the one hand, upregulation of this clearance pathway can reduce the accumulation of amyloid plaques and rescue neurodegeneration in various systems^[7-9]. On the other hand, $A\beta$ may also be generated in autophagosomes, which appear to contain both the precursor [the amyloid precursor protein (APP)] and the enzyme Presenilin-1 responsible for the cleavage of APP to $A\beta$ ^[7,10]. Furthermore, autophagy is critical for the extracellular secretion of $A\beta$ peptides, as conditional knockout of the key autophagy gene, *Atg7*, in APP transgenic mice carrying the Swedish mutation was shown to cause less extracellular $A\beta$ secretion and fewer extracellular $A\beta$ plaques. However, these events were associated with toxic intracellular aggregation of $A\beta$, which likely causes neurodegeneration, and together with amyloidosis, memory deficit^[11]. Besides $A\beta$, there is also a link between hyperphosphorylated tau and autophagy. Despite being substrates of the ubiquitin-proteasome system, tau proteins are delivered to the autophagosome-lysosomal system for degradation, as autophagy impairment leads to the formation of tau oligomers and accumulation of tau insoluble species^[12]. Indeed, autophagy-inducing drugs like verapamil or felodipine induce the clearance of tau in a zebrafish system^[13]. Furthermore, autophagy impacts tau phosphorylation: autophagy-deficient mice display hyperphosphorylated tau, while autophagy induction reduces tau phosphorylation^[14-16]. Interestingly, tau *per se* modulates the autophagy pathway, with tau accumulation disturbing the integrity of lysosome membranes and retarding autophagosome maturation by

inhibiting the activity of histone deacetylase 6^[15].

AD-associated mutations have been found to affect the three main steps of the autophagy pathways: autophagosome synthesis, trafficking, and maturation/degradation^[17,18].

The expression of Beclin-1, an essential participant in autophagosome biogenesis, is decreased in the affected brain regions of AD patients compared to healthy individuals, likely due to its increased proteolytic cleavage mediated by caspase 3, which is highly activated in AD brains^[19]. Since a decline in Beclin-1 activity leads to a decrease in autophagosome synthesis, this is associated with ageing and neurodegeneration^[20].

Genome-wide association studies in AD identified variants of phosphatidylinositol binding clathrin assembly protein (PICALM), which also shows accelerated cleavage and, consequently, loss-of-function, in AD^[21,22]. PICALM is a clathrin adaptor protein, whose loss-of-function impairs endocytosis of VAMP2 and VAMP3 (soluble NSF attachment protein receptors - SNAREs involved in the fusion of autophagosome precursors), as well as VAMP8 (involved in the fusion of autophagosomes with lysosomes/late endosomes). Thus, PICALM depletion in AD brains inhibits autophagy at multiple steps: from autophagosome biogenesis to maturation and fusion steps, ultimately leading to tau accumulation^[17,23].

Studies based on mouse and cell models of AD have reported that the trafficking of autophagy-related compartments is selectively defective^[24]. Inhibiting the microtubule-mediated retrograde transportation of autophagosomes towards lysosomes (from the distal axonal regions towards the soma) leads to the rapid accumulation of immature autophagosomes in neurites, a morphology similar to that observed in the AD brains^[7]. Tau is known to be critical for this retrograde vesicle trafficking as it regulates the assembly and stabilization of neuronal microtubules^[25]. Importantly, tau gains a toxic function in AD models, generating neurofibrillary tangles, which may negatively impact neuronal vesicle trafficking^[26]. Reduced expression of the chaperonin CCT/TRiC complex, an important chaperone that assists the proper folding of tau, was found in the affected brain regions of AD patients^[27,28]. Apart from its direct action on tau, CCT is essential for maintaining the cytoskeleton structure in neurons and consequently to ensure proper trafficking of autophagosomes and their ultimate fusion with lysosomes. CCT impairment, by blocking autophagosome delivery/degradation, causes the accumulation of long-lived proteins and autophagy substrates, such as p62^[27].

In addition to the aforementioned factors regulating autophagic flux (PICALM and CCT), mutations of Presenilin-1 (PS1), known to cause early-onset familial Alzheimer's disease, impair autophagosome maturation/degradation via dysfunction of the lysosomal V-ATPase in patient-derived cells^[29]. Besides its role in the APP cleavage to A β as part of the γ -secretase complex, PS1 regulates N-glycosylation of the V-ATPase subunit V_oA1 essential for lysosomal acidification and subsequent degradation of autophagic content. Abnormally increased levels of acid sphingomyelinase (SMase) in AD brains lower TFEB levels by promoting its proteolysis^[30]. Conversely, chemical inhibition of this SMase with amitriptyline increased the degradation of p62 and LC3-II, and delayed the disease development in APP/PS1 mouse models of AD^[17,30]. Additional genetic risk factors for AD, such as the *APOE4* variant, mutant APP or APP duplication, increase Rab5-mediated endocytosis and cause non-degraded lipids and proteins to accumulate in swollen lysosomes. Indeed, AD mice models show lysosomal dysfunction with subsequent accumulation of autophagy substrates^[7,10].

Autophagy modulation as therapy in AD

Autophagy induction may be a promising therapeutic strategy in AD [Table 2]. Autophagy stimulation can be achieved either chemically [using mechanistic target of rapamycin (mTOR)-dependent or -independent small molecule autophagy enhancers] or through gene therapy approaches, as summarised below^[6,31].

The first group of chemical compounds, which inhibit the negative autophagy regulator mTOR, such as rapamycin and other rapalogs (eg. temsirolimus), reduce the levels of hyperphosphorylated tau and A β , and consequently ameliorate the cognitive and memory deficits in multiple AD model mice (such as P301S mutant tau and PDAPP transgenic mouse models^[32,33]), likely due (at least in part) to autophagy activation in neurons^[34]. Rapamycin also reduces tau toxicity and increases the life-span of *Drosophila* (*Drosophila Melanogaster*) expressing wildtype or mutant R406W tau, as a result of enhanced autophagic clearance of soluble and insoluble tau^[35].

The second group of compounds mainly comprises the class of small molecule enhancers of autophagy that often signal through AMPK, which directly phosphorylates the AMPK substrate ULK1 and thereby induces autophagy^[36]. These molecules may be either direct AMPK modulators (such as metformin, bosutinib, nilotinib) or indirect upstream activators acting by lowering mitochondrial ATP production and/or intracytoplasmic Ca²⁺ levels^[37,38]. The direct AMPK modulators show beneficial effects in multiple transgenic APP mouse models^[39,40] and prevent the degeneration of neurons producing dopamine, a neurotransmitter involved in the regulation of cognitive and non-cognitive functions in a *Tg2576* AD mouse model^[41]. Thus, several of these AMPK-stimulating compounds have progressed to pilot clinical studies for AD, where metformin has shown promising early results with improved memory and executive function^[42,43].

The second group of compounds comprises a continuously growing list of molecules. Among them, the disaccharide trehalose activates AMPK by lowering ATP production by inhibiting GLUT transporters, and was shown to enhance the clearance of tau aggregates^[44-46]. The clearance of autophagy substrates can also be upregulated by a class of mood-stabilizing drugs, including lithium^[45,47]. Lithium administration results in lowering of the Ca²⁺ levels in the cytoplasm by reducing the intracellular levels of IP₃, decreasing the mitochondrial uptake of Ca²⁺ and consequently ATP production, causing AMPK activation^[48]. Lithium additionally upregulates autophagy by inhibiting GSK3 β , reducing the phosphorylation of tau and the formation of A β plaques in pre-pathological AD mouse models (APP^{swe}/PS1A246E)^[49]. In clinical trials, lithium treatment showed reduced cognitive decline over a two year period compared to the placebo group, accompanied by enhanced A β clearance from the brain^[50]. Other molecules that likely signal through the IP₃-Ca²⁺-AMPK axis include the class of imidazoline receptor agonists, such as clonidine and rilmenidine^[51]. Treatment with these two drugs increase the clearance of mutant tau and ameliorate the motor and morphological defects in an experimental zebrafish model expressing the human A152T disease-associated tau variant^[52]. Another class of compounds, the L-type Ca²⁺ channel antagonists (felodipine and verapamil) block the influx of extracellular Ca²⁺ and likely activate autophagy via the aforementioned Ca²⁺-AMPK axis. Consequently, apart from their antihypertensive effects, these drugs improve the clearance of insoluble tau and ameliorate the morphological defects seen in two transgenic zebrafish tauopathy models^[53]. Elevated intracytosolic Ca²⁺ levels activate calpains, which inhibit autophagy. The calpain inhibitor A-705253 improves clearance of hyperphosphorylated tau and A β and has beneficial cognitive effects in a triple transgenic AD mouse model (bearing disease-associated mutations in three genes: PS1M146V, APP^{swe}, TauP301L)^[54].

While the described chemical compounds that upregulate autophagy offer clear beneficial effects in multiple AD *in vivo* models, it is important to acknowledge that many of these compounds may also exert beneficial

Table 2. Therapeutic strategies to modulate autophagy and their effects on disease models (see main text for references)

Mechanisms of drugs	Examples
Rapamycin: mTOR inhibition	AD: Rapamycin and other rapalogs reduce the levels of hyperphosphorylated tau and A β , and improve cognitive and memory deficits in AD mice and <i>Drosophila</i> PD: Rapamycin treatment increases the clearance of mutant α -syn and attenuates toxicity in PC12 cells, dopaminergic neurons and PD animal models HD: Rapamycin and its derivatives facilitate clearance of PolyQ aggregates and reduce toxicity in HD <i>Drosophila</i> and mice ALS: Rapamycin treatment decreases TDP-43 inclusions and forebrain neurodegeneration in a <i>FTD</i> mouse model. Rapamycin also reduces neuronal TBPH aggregates in an ALS <i>Drosophila</i> model, as well as partially rescuing lifespan and locomotor defects in these animals Ageing: Rapamycin increases mortality in mice and life extension in <i>Drosophila</i>
SMERs: mTOR-independent autophagy	PD: SMERs induce autophagic clearance of A53T α -syn <i>in vitro</i> independently of mTOR HD: SMERs trigger the elimination of mHTT aggregates and ameliorate toxicity in cell culture and in HD <i>Drosophila</i> models
Metformin, bosutinib, nilotinib: AMPK activation	AD: Metformin, bosutinib and nilotinib prevent the degeneration of dopaminergic neurons, the loss of which affects cognitive and non-cognitive functions in a Tg2576 AD mice PD: Metformin reduces levels of phosphorylated α -syn Ser129 and rescues mitochondrial dysfunction in SH-SY5Y and <i>Drosophila</i> models Nilotinib partially rescues disease phenotypes in both A53T and PD mice and improves cognitive and motor skills in PD patients Resveratrol induces α -syn clearance, possibly via interaction with SIRT1 in SH-SY5Y and PC12 cells HD: Metformin significantly clears PolyQ aggregates and improves motor and behavioural dysfunction in PolyQ disease mice
Trehalose: AMPK activation	AD: Trehalose induces tau clearance and neuronal survival in AD <i>tauP301S</i> mice ALS: Trehalose ameliorates early motor phenotypes in the G93A SOD-1 ALS mice, but fails to delay end-stage motor phenotypes HD: Trehalose enhances the degradation of mHTT aggregates in HD mice
Lithium, Clonidine, rilmenidine: IP ₃ -Ca ²⁺ -AMPK axis	AD: Lithium reduces tau phosphorylation and the formation of A β plaques in pre-pathological AD mice. Clonidine and rilmenidine induce mutant tau clearance and ameliorate motor defect in A152T tau zebrafish PD: Lithium increases the clearance of A53T and A30P mutant α -syn in inducible PC12 cell models and ameliorates phenotypes associated with PD in aged Parkin mutant tg mice HD: Clonidine enhances removal of aberrant mHTT aggregates in HD in an mTOR-independent manner
Felodipine, verapamil: Ca ²⁺ -AMPK axis	AD: Felodipine and verapamil improve the clearance of insoluble tau and ameliorate the morphological defects seen in two transgenic zebrafish tauopathy models PD: Felodipine decreases the levels of A53T α -syn in a PD mouse model and ameliorates neurodegeneration HD: Felodipine decreases mHTT in a HD mouse model and ameliorates signs of disease
A-705253: Calpain inhibition	AD: A-705253 improves clearance of hyperphosphorylated tau and A β and has beneficial cognitive effects in 3xtg AD mice HD: Downregulation of Calpain decreases mHTT aggregates and is neuroprotective in HD mice and <i>Drosophila</i>

effects via autophagy-independent mechanisms. Thus, targeting key autophagy genes through genetic approaches may enable more specific therapeutic strategies.

One of the most common strategies to genetically stimulate autophagy is by lentiviral-mediated overexpression of *BECN1*, which encodes Beclin-1^[19]. This strategy ameliorates the amyloid pathology in the hippocampus and cortex of APP transgenic mice. Beclin-1-mediated autophagy has also been successfully induced by lentiviral Parkin transduction and was shown to clear intracellular A β in a triple transgenic AD mouse model (mutations in three genes: *PS1M146V*, *APPswe*, *TauP301L*)^[55]. Another elegant approach, using a knock-in point mutation in *BECN1*(F121A) that blocks the interaction with its canonical inhibitor Bcl-2, causes a significant reduction in A β levels and increases the life-span of *5xFAD* mice (expressing a combination of 5 familial AD mutations in human *PS1* and *APP* genes)^[56]. Similar to *BECN1* overexpression, ATG5-induced autophagy ameliorates the morphological defects and enhances the tau clearance in an A152T tau zebrafish model^[52].

AD is a complex disease that results in impairment at multiple steps in the autophagy pathway, from autophagosome biogenesis to lysosomal functioning. In the AD scenarios with compromised autophagosome degradation within lysosomes, overexpression of *BECN1* and *ATG5* genes mainly induces the formation of autophagosomes, with little effect on lysosomal biogenesis, and has the theoretical potential to lead to the accumulation of nondegraded autophagosomes, which may further worsen pathology (depending on the extent of defective autophagosome clearance). To improve lysosomal function, studies using genetic ablation of cystatin B (an endogenous inhibitor of lysosomal cysteine proteases) successfully alleviated the memory deficits and amyloid pathology in a *TgCRND8* mouse model of aggressive AD amyloidosis^[57]. Thus, combined approaches that act to enhance overall autophagy flux, by simultaneously upregulating the autophagosome synthesis and improving the lysosomal functioning, would be ideal therapeutic strategies. Therefore, an interesting target for drug development may be TFEB, as this master transcription factor controls the synthesis of both autophagosomes and lysosomes^[38,58]. Indeed, TFEB overexpression improves the behavioural defects and enhances the clearance of hyperphosphorylated tau in a *rTg4510* mouse model of tauopathy^[59].

Parkinson's disease

PD is the most common neurodegenerative movement disorder, primarily associated with progressive loss of motor control, often accompanied by cognitive decline. It is characterized by the presence of intraneuronal inclusions known as Lewy bodies and Lewy neurites enriched with filamentous forms of α -synuclein (α -syn)^[60], which is encoded by *SNCA*. Multiplications of the *SNCA* locus cause autosomal dominant forms of PD. Levels of α -syn correlate with disease severity^[61]. Defective autophagy has been implicated as central in the aetiology and pathogenesis of PD. Similar to AD, mutations causing PD appear to affect different stages of the autophagy itinerary, such as autophagosome biogenesis or fusion with lysosomes^[62].

The presence of inclusion bodies containing α -syn affects autophagosome maturation and fusion with lysosomes, resulting in decreased protein degradation^[63], suggesting that the presence of α -syn compromises autophagic flux. Overexpression of α -syn *in vitro* and *in vivo* results in compromised autophagosome biogenesis through the inhibition of Rab1, causing mislocalisation of ATG9^[64]. A mutation in *VPS35*, which encodes a component of the retromer complex, results in an autosomal dominant form of PD, impairs autophagy, causes ATG9 mislocalisation and inhibits autophagosome formation^[65]. A recent study in an α -synucleinopathy *Drosophila* model suggests that α -syn expression impairs autophagic flux in ageing adult neurons by disrupting the F-actin cytoskeleton^[66].

There is increasing evidence of lysosomal defects in PD. Lysosomal impairment has been shown to cause accumulation of α -syn aggregates in the nervous system^[67]. Autosomal dominant mutations in the *LRRK2* gene appear to affect endosome-to-lysosome trafficking. Overexpression of mutant *lrrk*, analogous to the most common PD causing mutation in human *LRRK2* (G2019S), led to defective lysosomal positioning mediated by Rab7 and enlarged lysosomes containing nondegraded contents^[68].

Heterozygous mutations in the gene encoding the lysosomal enzyme glucocerebrosidase (GBA) are the most commonly known genetic risk factors for PD. Induced pluripotent stem cells-derived neurons from PD patients with *GBA* mutations showed increased α -syn levels, as well as autophagic and lysosomal defects^[69].

Furthermore, the PD-associated genes *ATP13A2* and *SYT11* regulate autophagy through a pathway mediated by TFEB^[70]. Depletion of *ATP13A2* *in vitro* has been shown to decrease levels of SYT11. The

decrease in SYT11 can account for the lysosomal dysfunction and impaired autophagosome degradation resulting from ATP13A2 deficiency, since SYT11 overexpression in ATP13A2 knockdown cells was able to rescue the autophagy defects in these cells^[70].

Autosomal recessive mutations in *PINK1* or *Parkin*, two genes encoding proteins that are key regulators of a form of mitophagy, cause early onset Parkinsonism^[71]. While these genes have led to suggestions that impaired mitochondrial quality control may contribute to PD, we have not focussed on this domain in this review focussing on ageing neurodegenerative diseases, as these Mendelian mutations manifest with disease in younger patients.

Autophagy modulation as therapy in PD

Inhibition of mTOR with rapamycin treatment increases the clearance of WT and PD-related mutant forms (A30P and A53T) of α -syn in PC12 cells^[72], thereby attenuating toxicity in both dopaminergic neurons^[73] and animal models of PD [Table 2]^[74,75].

Various mTOR-independent autophagy inducers have shown beneficial effects in PD models. Several novel small-molecule enhancers of rapamycin (SMERs) - SMER10, SMER18, and SMER28 - were found to induce autophagic clearance of A53T α -syn *in vitro* independently of mTOR^[76].

Another group of compounds that have shown beneficial effects in PD are AMPK activators. Metformin, a drug currently used in the treatment of type 2 diabetes, reduces levels of Ser-129 phosphorylated α -syn in SH-SY5Y cells expressing α -syn^[77] and partially rescues the mitochondrial dysfunction in genetic *Drosophila* models of PD^[78]. Nilotinib, a tyrosine kinase inhibitor, induces autophagy, and partially rescues disease phenotypes in both A53T and lentiviral α -syn overexpression PD mouse models by inhibiting phosphorylation of BCR-ABL^[79]. Clinical trials of nilotinib in PD patients showed improved cognitive and motor skills^[80]. Resveratrol, a natural plant phenol, induces AMPK-dependent autophagy and α -syn clearance in SH-SY5Y and PC12 cells, possibly via SIRT1, a histone/protein deacetylase that activates autophagy via deacetylation of autophagy proteins^[81-83]. The mTOR-independent autophagy inducer lithium increases the clearance of A53T and A30P mutant α -syn in inducible PC12 cell models^[47] and reduces phenotypes associated with PD in an aged Parkin mutant transgenic mouse model^[84]. In addition to lithium, other mood-stabilizing drugs, such as sodium valproate and carbamazepine, have been shown to induce autophagy in SH-SY5Y cells exposed to rotenone-induced toxicity^[85]. Upon treatment with trehalose, autophagy was induced and enhanced clearance of α -syn both *in vitro*^[86] and *in vivo*^[87]. Additionally, felodipine decreases the levels of A53T α -syn in a mouse model of PD and ameliorates neurodegeneration^[13].

Huntington's disease

HD is a heritable progressive neurodegenerative disease categorized as a polyglutamine (PolyQ) disease, along with other diseases such, dentatorubralpallidoluysian atrophy, spinal and bulbar muscular atrophy (SBMA), spinocerebellar ataxias (SCA) types 1, 2, 6, 7, 17, as well as Machado-Joseph disease (MJD/SCA3)^[88]. Most PolyQ diseases are autosomal dominant except for SBMA, which is an X-linked disorder characterised by neurological, psychiatric and motor symptoms^[89].

Each of these PolyQ diseases are caused by abnormal CAG repeat expansions that encode a PolyQ tract within specific genes. The lengths of PolyQ tracts vary in the different PolyQ diseases and cause formation of toxic oligomers and aggregates. Moreover, the length of the PolyQ tracts negatively correlates with the age of disease onset^[90]. PolyQ diseases have been associated with dysfunctional autophagy and an altered

endo-lysosomal network required for the elimination of aggregates^[91].

HD, the most common PolyQ disease, results from an abnormal CAG repeat expansion (> 35 repeats) encoding a PolyQ tract at the N-terminus of the huntingtin (*Htt*) gene^[92]. PolyQ expansions in mutant huntingtin (mHTT) promote the accumulation of perinuclear cytoplasmic aggregates and intranuclear inclusion bodies in neurons^[93,94,95]. Defects have been observed at different stages of the autophagy itinerary in Huntington's disease. The non-aggregated mutant huntingtin impacts starvation-induced autophagosome biogenesis by impairing the ability of wildtype ataxin-3 to deubiquitinate Beclin-1 and protect this protein from proteasomal degradation^[96]. HD is also associated with defective autophagic cargo recognition^[97]. Wildtype HTT acts as a scaffold protein which interacts with autophagy components to facilitate the degradation of selective autophagy cargos^[98,99]. Defects in cargo recognition caused by mHTT inhibit organelle sequestration through autophagy and consequently result in abnormal intracellular lipid stores and altered mitochondria in mHTT mice^[97]. HTT plays an important role in the regulation for axonal transport of autophagosomes via huntingtin-associated protein-1 (HAP1). In particular, loss of HTT or HAP1 disrupts the retrograde axonal transport of autophagosomes^[100]. Retrograde transport is controlled by HTT and the microtubule motor protein dynein. Suppression of dynein inhibits autophagosome-lysosome fusion and aggravates the phenotype in HD *Drosophila* and mice^[101]. Moreover, phosphorylation of HTT by AKT controls anterograde transport via kinesin1 recruitment^[102]. Indeed, impaired autophagosome dynamics by mHTT expression also results in the inefficient degradation of dysfunctional mitochondria and mHTT aggregates in HD neurons, indicating that HTT controls its own elimination^[100,101]. In addition, mHTT is involved in the regulation of post-Golgi trafficking. mHTT reduces the interaction with the optineurin/Rab8 complex and subsequently perturbs post-Golgi trafficking to lysosomes, which could result in dysfunctional autophagosome/lysosome dynamics^[103]. A striatal-specific protein, Rhes, directly binds mHtt and accelerates mHTT-mediated cytotoxicity^[104]. In the striatum, Rhes binds to Beclin-1 and promotes the dissociation of Bcl-2 and Beclin-1, leading to autophagy activation. However, mHTT expression interferes with Rhes-mediated autophagy induction^[104,105]. Lower levels of Rhes are observed in HD, whereas overexpression of Rhes ameliorates HD-associated phenotypes in mice^[105]. The V271A polymorphism in the core autophagy gene *ATG7* is associated with earlier onset of HD, suggesting that *ATG7* may be implicated in HD, although the exact mechanisms associated with this polymorphisms have yet to be fully elucidated^[106]. In addition, PolyQ tracts in mHTT impair mitophagy and have been associated with increased oxidative stress and the accumulation of damaged mitochondria^[107]. Interestingly, compensatory upregulation of CMA is observed in the early stages of HD, possibly as response to macroautophagic dysfunction^[108].

Autophagy modulation as therapy in HD

Pharmacological and genetic approaches to upregulate autophagy in PolyQ diseases have shown that increased clearance of PolyQ aggregates via autophagy can ameliorate PolyQ disease pathology [Table 2]^[91]. In contrast, inhibition of autophagy has been shown to cause cytotoxicity as well as increased polyQ aggregates accumulation *in vivo*^[109].

Depletion of mHTT itself in symptomatic HD mice significantly reduced aggregation in the brain. Deletion of the PolyQ tract within HTT (Δ Q-htt) induces mTOR-independent autophagy in neurons. Interestingly, expression of Δ Q-htt in PolyQ-HTT knock-in (Hdh140Q/+) mice has protective effects, decreasing HTT neuropil aggregates, mitigating motor and behavioural defects as well as promoting longevity^[110]. mTOR-dependent autophagy activation has been shown to have beneficial effects in PolyQ disease. Rapamycin and its derivatives reduced toxicity. In addition, the rapamycin analogue temsirolimus facilitated clearance of PolyQ aggregates mediated by autophagy activation and rescued pathology in HD *Drosophila* and mice^[109,111]. mTOR-independent autophagy activation via trehalose enhances the degradation of mHTT

aggregates. Combination treatment of rapamycin and trehalose synergistically accelerated clearance of mHTT aggregates in HD *Drosophila*^[86]. SMER10, 18, and SMER28, mTOR-independent autophagy inducers, trigger the elimination of mHTT aggregates and ameliorate toxicity in cell culture and in *Drosophila* HD models^[76]. Interestingly, dual treatment SMERs with rapamycin had enhanced protective effects. Lithium treatment decreased soluble and aggregated mHTT, correlating with decreased toxicity^[112]. Induction of autophagy in both an mTOR-independent and -dependent manner using a combination therapy of lithium and rapamycin leads to protection against neurodegeneration in an HD *Drosophila* model^[112]. The G_i signalling activator clonidine, which induces mTOR-independent autophagy and regulates cAMP or IP₃, enhances removal of aberrant mHTT aggregates in HD^[51]. Other autophagy inducers such as berberine, rilmenidine, and metformin significantly clear PolyQ aggregates and improved motor and behavioural dysfunction in PolyQ disease mice^[113-115].

Downregulation of calpain induces autophagy, decreasing the number of mHTT aggregates and consequently has neuroprotective effects in HD mice and *Drosophila*. Although calpain inhibition is neuroprotective in HD mouse models, prolonged brain overexpression of calpastatin, a calpain inhibitor, does not cause obvious deleterious effects in mice^[116]. Depletion of the autophagy adaptor protein Alfy exacerbates mutant huntingtin toxicity in mice and HD patient-derived neuronal model^[117]. Conversely, overexpression of Alfy induces selective degradation of aggregated proteins as well as neuroprotection in neuronal and *Drosophila* HD models, suggesting that Alfy-mediated selective autophagy alleviates HD neuropathology^[117,118]. Thus, improved cargo recognition by autophagy adaptors may be a potential target for therapeutic strategy.

Expression of TFEB, which promotes lysosome biogenesis and autophagy, leads to increased lysosomal activity in the striatum of HD mice, resulting in decreased levels of mHTT^[119]. In addition, a previous study reported that both macroautophagy and CMA function in the degradation of the Htt fragment containing amino acids 1-552. Interactions between mHTT and the CMA components heat shock protein cognate 70 and lysosome-associated protein 2A (LAMP-2A) enable the uptake and lysosomal degradation of Htt-552, whereas inhibition of the CMA pathway suppresses Htt-552 degradation^[120]. Acetylation of mHTT by CBP activation or HDAC1 inhibition triggered mHTT degradation and enhanced neuroprotection through the autophagic-lysosomal pathway in primary neurons and *Caenorhabditis elegans* (*C. elegans*) HD model^[121,122].

ALS/FTD

In recent years, the neurodegenerative diseases ALS and FTD have become widely accepted as divergent presentations of a common neurodegenerative disease process^[123]. ALS is characterised by loss of lower motor neurons from the brainstem/spinal cord and upper motor neurons from the motor cortex, while FTD involves loss of neurons from the prefrontal and temporal cortices. However, many patients with ALS exhibit symptoms traditionally associated with FTD (cognitive impairment and behavioural changes) and patients with FTD can develop motor neuron involvement resulting in progressive muscle weakness^[123].

ALS/FTD are age-related diseases, typically manifesting in the fifth or sixth decade of life^[124]. Conversely, ALS transgenic mouse models are reported to exhibit abnormalities in axonal transport, sensorimotor development and neuronal excitability as early as the neonatal period, despite not exhibiting neurodegenerative phenotypes until mature adulthood^[125-127]. This observation has prompted speculation that normal ageing is a prerequisite for the ALS/FTD neurodegenerative disease process^[124].

One possible explanation is that normal age-related phenotypes exhibited by relevant cell types increase vulnerability to ALS/FTD pathology^[128]. Relevant to ALS, various motor neuron deficits have been

associated with normal ageing. Motor neuron number decreases in older rats (22 months old compared with 6 months old)^[129] and humans (diminished motor neuron population evident over 60 years old)^[130]. Energy homeostasis is also impaired, with motor neurons from aged mice demonstrating less efficient electrophysiological properties (20 months old compared with 12 months old)^[131] and motor neurons from elderly human donors (68-99 years old) exhibiting mitochondrial dysfunction^[132]. These age-related phenotypes and others are likely to lower the susceptibility threshold for ALS/FTD-specific neurodegeneration.

Reduced autophagy in skeletal muscle is another factor liable to increase vulnerability to ALS pathology as part of normal ageing. Central to less “neuron centric” models of ALS pathology is the notion that skeletal muscle is structurally and functionally interconnected with lower motor neurons. Decreased lysosomal activity leading to impaired autophagic flux and lower expression of core autophagy proteins (ATG7 and LC3) is reported in skeletal muscle from older mice^[133,134], with ATG7 and LC3 expression also decreased in skeletal muscle from sedentary elderly human donors^[134].

Around 5%-10% of ALS and 30%-50% of FTD cases are familial, with mutations in more than 25 genes associated with ALS/FTD^[135]. Sporadic and familial ALS/FTD are clinically identical. Models based on disease-causing mutations have implicated dysfunction at several points in the autophagy pathway in ALS/FTD pathogenesis. For instance, ALS-linked mutations in the autophagy receptor optineurin (Q398X and E478G) are reported to block autophagosome-lysosome fusion by disrupting myosin VI-mediated lysosomal trafficking in motor neuron-like NSC-34 cells^[136]. Phosphorylation by the protein kinase TBK1 has been shown to improve the efficiency of optineurin as an autophagy receptor, thereby increasing autophagy substrate clearance^[137]. ALS/FTD-causing mutations identified in TBK1 that impair binding and phosphorylation of optineurin^[138,139] are therefore expected to reduce autophagy substrate recognition and clearance. Numerous ALS/FTD-linked mutations have also been identified in the autophagy receptor p62 (also known as SQSTM1)^[140-142], with ALS-associated L341V p62 exhibiting defective binding to the core autophagy protein LC3 and impaired recruitment to autophagic vesicles in NSC-34 cells^[143].

Other ALS/FTD-causing mutations linked to autophagy include a splice-site mutation in the ESCRT (endosomal sorting complex required for transport) protein CHMP2B, which produces a truncated final protein^[144,145]. Truncated CHMP2B blocks flux through the autophagy pathway in cultured primary neurons from rat cortex and photoreceptor neurons in fly eye sections, resulting in autophagosome accumulation^[145]. The ATPase p97 (also known as VCP) also exhibits an ALS/FTD-linked mutation (R155H) that disrupts clearance of damaged lysosomes by autophagy. MEFs and muscle fibres from human donors expressing R155H p97 consequently accumulate defective lysosomes^[146]. R155H p97 and A232E p97 (another disease-related mutation) additionally drive accumulation of stress granules (nontranslating messenger ribonucleoprotein aggregates) in HeLa cells, suspected to result from defective autophagy^[147]. Very recently, p97 has been shown to function in early autophagy initiation through the core autophagy protein Beclin-1^[148]. Models based on disease-causing mutations therefore indicate defects in autophagy substrate recognition, autophagosome biogenesis and autophagosome-lysosome fusion, depending on the functional effects of the mutations on VCP activities.

Cytosolic protein aggregates, which are autophagy substrates, are found almost universally in ALS/FTD and typically comprise TDP-43 (TAR DNA binding protein 43), SOD-1 (superoxide dismutase 1), FUS (fused in sarcoma) and DPR (dipeptide repeat) proteins in various combinations^[149]. Tau, p62 and ubiquitin can also feature in these cytosolic inclusions. Consequently, further experimental evidence linking autophagy to ALS/FTD stems from autophagic degradation of ALS/FTD-associated aggregate-prone proteins.

A hexanucleotide (GGGGCC) repeat expansion in an untranslated region of the C9ORF72 transcript is the most common genetic abnormality identified in familial and sporadic ALS/FTD^[150-152]. Hexanucleotide repeat expanded C9ORF72 generates DPR proteins, which adopt aggregate-prone conformations and form cytosolic inclusions^[153]. Work in motor neuron-like NSC-34 cells indicates DPRs are predominantly cleared by autophagy^[154]. Loss of wildtype C9ORF72 function has also been identified in ALS/FTD caused by hexanucleotide repeat expanded C9ORF72^[155]. A “double-hit” pathogenic mechanism has been proposed, given that wildtype C9ORF72 is suggested to participate in both autophagosome biogenesis and lysosome-dependent autophagic flux^[149], whereby the hexanucleotide repeat expansion in C9ORF72 both generates DPR proteins and promotes neurotoxic DPR protein accumulation by impairing autophagy^[156].

Cytosolic aggregates of the RNA and DNA binding protein TDP-43 are found in almost all patients ALS, as well as over 40% of patients with FTD^[157]. While soluble (monomeric) TDP-43 is predominantly degraded by the ubiquitin-proteasome system, clearance of aggregated (oligomeric) TDP-43 requires autophagy^[158,159]. Since TDP-43 depletion has been shown to downregulate autophagy by decreasing expression of the core autophagy protein ATG7^[160], pathogenic feedback is again possible due to TDP-43 sequestration into aggregates causing loss-of-function.

Regarding other aggregate-prone proteins associated with ALS/FTD, autophagy is reported to degrade soluble and aggregated SOD-1, thereby preserving viability of mouse neuroblastoma cells expressing ALS/FTD-associated SOD-1 mutants^[161]. FUS cytosolic inclusions are commonly decorated with the autophagy receptor p62^[162]. However, interactions between autophagy and FUS-mediated ALS/FTD pathology are not well characterised. Finally, mitophagy (mitochondrial-selective autophagy) is known to clear defective mitochondria, which accumulate in ALS/FTD and compromise neuronal health^[163].

Autophagy modulation as therapy in ALS/FTD

There are currently only two FDA-approved drugs for ALS (riluzole and edavarone), which have limited efficacy in slowing disease progression, and no FDA-approved drugs for FTD^[164]. Autophagy modulation is therefore, an attractive therapeutic avenue in ALS/FTD [Table 2]. However, since defects in both autophagosome biogenesis and autophagic flux have been identified in ALS/FTD, autophagy inducers are clearly not panaceas for many forms of ALS/FTD.

A phase 2 clinical trial of rapamycin in ALS (RAP-ALS trial) has recently finished^[165], with the results now awaited. This trial follows on from promising animal studies. For example, rapamycin treatment decreased TDP-43 inclusions and forebrain neurodegeneration in a FTD mouse model^[166]. Increased autophagy was observed in the mouse forebrain after rapamycin treatment, with motor and behavioural phenotypes also ameliorated^[166]. Rapamycin also reduced neuronal TBPH (TAR DNA binding protein 43 homolog) aggregates in a *Drosophila* ALS model, as well as partially rescuing lifespan and locomotive defects in these animals^[167]. On the other hand, when studied using ALS/FTD models not based on TDP-43, rapamycin exhibited less therapeutic potential and sometimes worsened neurodegenerative phenotypes. Rapamycin accelerated motor neuron degeneration and shortened lifespan in a transgenic mouse model of ALS expressing disease-causing mutant SOD-1 (human G93A SOD-1), for instance^[168]. This is suggested to relate to impaired autophagic flux in these animals, which is exacerbated by autophagy induction using rapamycin^[168].

Experimental evidence concerning mTOR-independent autophagy inducers in ALS/FTD is similarly mixed. Treatment with trehalose ameliorated early motor phenotypes in the G93A SOD-1 transgenic mouse model of ALS, but failed to delay end-stage motor phenotypes and did not extend survival. This was despite

increased autophagy and reduced SOD-1 levels in the spinal cords of these animals after trehalose treatment^[169]. Lithium showed initial therapeutic promise in patients with ALS^[170]. However, subsequent higher powered clinical trials reported no benefit from lithium treatment on either disease progression or survival in patients with ALS^[171-174]. The heterogeneous nature of ALS/FTD pathology, which extends to defects at multiple points in the autophagy pathway, therefore makes unlocking the therapeutic potential of autophagy modulation in ALS/FTD complicated, necessitating further study.

AUTOPHAGY AND AGEING

While the accumulation of protein aggregates is traditionally associated with neurodegenerative diseases, studies of post-mortem brains of aged individuals, who were not diagnosed with neurological conditions, shows the presence of amyloid plaques, neurofibrillary tangles, Lewy bodies, synaptic dystrophy and neuronal loss, consistent with the notion that the integrities of degradative pathways are challenged with age^[175]. Indeed, there is increasing evidence suggesting that autophagy declines during ageing in many organisms and that this reduction plays a role in the functional deterioration of biological functions with age^[176].

In multiple studies, autophagy gene transcripts decrease with age in the brain and muscle of *Drosophila*^[177-179]. Transcriptional downregulation of *ATG5*, *ATG7* and *BECN1* with age was found in post-mortem human brains^[180]. This correlates with the age-related decrease in autophagy proteins in the mouse hypothalamus^[181] and human muscle^[134], as well as lysosomal proteins in rat livers^[182]. In a spatiotemporal analysis of autophagy in *C. elegans*, an age-dependent decrease in the numbers of autolysosomes and autophagosomes was observed in the intestines, muscles and neurons^[183], corresponding with another study showing decreased autophagic activity in whole-body extracts of aged *C. elegans*^[184]. Electron microscopy analyses have shown an accumulation of autophagic vacuoles and decreased ability to clear autophagic vesicles in mouse and rat livers with age^[185-187].

Additionally, studies have shown a correlation between autophagy and lifespan. In *C. elegans*, decreased expression of orthologues of the mammalian *ATG1*, *ATG7*, *ATG12*, *BECN1* and *ATG18* lead to shortened lifespan^[188]. Deletion of core autophagy genes, such as *Atg7* in flies, reduce life-span and cause accumulation of aggregated proteins in degenerating neurons^[189], which was supported by findings revealing reduced lifespan in mutant flies with reduced *Atg1* and *Atg8* expression^[179]. A study in senescence-accelerated mouse-prone 8 mice, a rodent model with accelerated ageing, showed an accumulation of autophagic vesicles in hippocampal neurons along with deficits in learning and memory with increasing age^[190]. The age-dependent decline in autophagic function and lysosomal degradation was prevented with dietary restriction^[191,192].

Overexpression of specific autophagy genes extends lifespan in flies and mice^[179,193,194]. Additionally, overexpression of the TFEB orthologue, helix-loop-helix transcription factor *hlh-30* in *C. elegans* extended lifespan in an autophagy-dependent manner^[195]. Disruption of the interaction of Beclin-1 with its negative regulator Bcl-2 achieved by introducing a point mutation in *BECN1(F121A)* in mice led to increased lifespan and decreased age-related renal and cardiac pathological changes and spontaneous tumorigenesis^[196].

In addition to bulk autophagy, a decline in selective forms of autophagy, such as the autophagic degradation of mitochondria has been observed in aged worms, flies, mice and humans^[37,197], possibly accounting for the increased presence of inefficient and toxic mitochondria that have been implicated in neurodegenerative and inflammatory pathologies^[198]. In accordance with this notion, upregulation of mitophagy, as well as

pharmacological induction of mitophagy was found to improve longevity in worms^[199-201].

It is evident that the progressive loss of the degradative capacity can lead to an accumulation of toxic proteins, as autophagy gradually decreases with age. As mentioned earlier, autophagic decline is especially relevant to neurons, as post-mitotic cells are unable to segregate dysfunctional proteins and organelles from daughter cells using mitosis, resulting in an increased reliance on autophagy. This can have a detrimental impact on neuronal health and may play a role in manifestation of neurodegenerative diseases.

Therapeutic potential of autophagy in ageing

In addition to therapeutic potential in neurodegenerative conditions, autophagy upregulation is also suggested to ameliorate phenotypes associated with normal ageing and extend healthy lifespan. Caloric restriction drives autophagy-dependent lifespan extension in both *S. cerevisiae* (budding yeast)^[202,203] and *C. elegans*^[204]. Similar results are reported using *daf-2* mutant *C. elegans*, which exhibit lifespan extension due to defective insulin-like signalling dependant on autophagy^[188,205]. Transgenic mice moderately overexpressing the core autophagy protein ATG5 in all tissues live 17% longer than wildtype controls and also exhibit “anti-ageing” phenotypes, such as enhanced insulin sensitivity^[194]. Whether these phenotypes result directly from increased autophagy is not addressed in this study. More recently however, the lifespan extensions and milder age-related heart and kidney phenotypes exhibited by mice expressing constitutively active Beclin-1 (F121A mutant) has been shown to require increased flux through the autophagy pathway^[196]. Pharmacological upregulation of autophagy has also been suggested to extend lifespan. For example, a multicentre trial has demonstrated that feeding aged (600 days old) mice rapamycin significantly increases age at 90% mortality^[206]. This result has also been seen in *Drosophila* with rapamycin-induced lifespan extension dependent on expression of the core autophagy protein ATG5^[207].

CONCLUSION

Although we can currently only speculate that autophagy induction would ameliorate age-related phenotypes in the brain and nervous system, neurodegeneration undeniably occurs predominantly against the backdrop of normal ageing. Accordingly, there is growing appreciation that research into normal ageing and age-related neurodegenerative diseases should be drawn closer together^[128]. This includes incorporating ageing into models of neurodegenerative disease, which should increase model fidelity, leading to more efficient discovery of translatable disease-modifying therapies. Other considerations relevant to autophagy modulation as therapy in age-related neurodegenerative diseases include the mechanism through which autophagy is upregulated. The mTOR pathway for example, regulates numerous autophagy-independent processes such as cell growth and immunity^[208]. Hence, autophagy inducers that function by inhibiting mTOR may have deleterious side effects. Variations in the primary disease-causing mechanisms exhibited by different individuals with the same neurodegenerative disease is another important consideration. Specifically, individuals with different variants of the same neurodegenerative disease may exhibit defects at different locations in the autophagy pathway. Another consideration is variations in pathology between cell types in the same individual at different stages of a neurodegenerative disease. It is also possible for autophagy defects to be of differential importance as a neurodegenerative disease progresses. Accordingly, an autophagy inducer might be beneficial in slowing the onset of a neurodegenerative condition, but not improve (or even worsen) end-stage symptoms. A nuanced approach that takes into consideration the complex relationship between autophagy, ageing and neurodegeneration is therefore required if the therapeutic potential of autophagy modulation is to be realised and produce new disease-modifying treatments for age-related neurodegenerative diseases.

DECLARATIONS

Authors' contributions

Contribute equally to the research and drafting of the article: Karabiyik C, Frake RA, Park SJ, Pavel M, Rubinsztein DC

Coordinate the first draft: Karabiyik C

Edite the article: Rubinsztein DC

Availability of data and materials

Not applicable.

Financial support and sponsorship

This work was supported by UK Dementia Research Institute (funded by the MRC, Alzheimer's Research UK and the Alzheimer's Society), Roger de Spoelberch Foundation, Alzheimer's Research UK, The Tau Consortium, Cambridge Centre for Parkinson-Plus, National Institute for Health Research Cambridge Biomedical Research Centre (Rubinsztein DC); Gates Cambridge Scholarship (Karabiyik C); a grant of the Romanian Ministry of Research, Innovation and Digitization, CNCS/CCCDI - UEFISCDI, project number PN-III-P1-1.1-PD-2019-0733, within PNCDI III (Pavel M); the South East Scotland Academic Foundation Programme (Frake RA). The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

Conflicts of interest

Rubinsztein DC is a consultant for Aladdin Healthcare Technologies SE, Drishti Discoveries, PAQ Therapeutics Inc. and Nido Biosciences. None of the other authors have any potential competing interests.

Ethical approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Copyright

© The Author(s) 2021.

REFERENCES

1. Cai Z, Zhao B, Li K, et al. Mammalian target of rapamycin: a valid therapeutic target through the autophagy pathway for Alzheimer's disease? *J Neurosci Res* 2012;90:1105-18. DOI PubMed
2. Cuervo AM. Autophagy and aging: keeping that old broom working. *Trends Genet* 2008;24:604-12. DOI PubMed PMC
3. Stolz A, Ernst A, Dikic I. Cargo recognition and trafficking in selective autophagy. *Nat Cell Biol* 2014;16:495-501. DOI PubMed
4. Tsukada M, Ohsumi Y. Isolation and characterization of autophagy-defective mutants of *Saccharomyces cerevisiae*. *Febs Lett* 1993;333:169-74. DOI PubMed
5. Rubinsztein DC, Bento CF, Deretic V. Therapeutic targeting of autophagy in neurodegenerative and infectious diseases. *J Exp Med* 2015;212:979-90. DOI PubMed PMC
6. Djajadikerta A, Keshri S, Pavel M, Prestil R, Ryan L, Rubinsztein DC. Autophagy Induction as a Therapeutic Strategy for Neurodegenerative Diseases. *J Mol Biol* 2020;432:2799-821. DOI PubMed
7. Boland B, Kumar A, Lee S, et al. Autophagy induction and autophagosome clearance in neurons: relationship to autophagic pathology in Alzheimer's disease. *J Neurosci* 2008;28:6926-37. DOI PubMed PMC
8. Tian Y, Bustos V, Flajolet M, Greengard P. A small-molecule enhancer of autophagy decreases levels of Abeta and APP-CTF via Atg5-dependent autophagy pathway. *FASEB J* 2011;25:1934-42. DOI PubMed PMC
9. Vingtxdeux V, Chandakkar P, Zhao H, d'Abramo C, Davies P, Marambaud P. Novel synthetic small-molecule activators of AMPK as enhancers of autophagy and amyloid- β peptide degradation. *FASEB J* 2011;25:219-31. DOI PubMed PMC
10. Nixon RA, Wegiel J, Kumar A, et al. Extensive involvement of autophagy in Alzheimer disease: an immuno-electron microscopy study. *J Neuropathol Exp Neurol* 2005;64:113-22. DOI PubMed
11. Nilsson P, Loganathan K, Sekiguchi M, et al. A β secretion and plaque formation depend on autophagy. *Cell Rep* 2013;5:61-9. DOI

[PubMed](#)

12. Li Q, Liu Y, Sun M. Autophagy and Alzheimer's Disease. *Cell Mol Neurobiol* 2017;37:377-88. [DOI PubMed](#)
13. Siddiqi FH, Menzies FM, Lopez A, et al. Author Correction: felodipine induces autophagy in mouse brains with pharmacokinetics amenable to repurposing. *Nat Commun* 2019;10:2530. [DOI PubMed PMC](#)
14. Caccamo A, Magri A, Medina DX, et al. mTOR regulates tau phosphorylation and degradation: implications for Alzheimer's disease and other tauopathies. *Ageing Cell* 2013;12:370-80. [DOI PubMed PMC](#)
15. Uddin MS, Mamun AA, Labu ZK, Hidalgo-Lanussa O, Barreto GE, Ashraf GM. Autophagic dysfunction in Alzheimer's disease: cellular and molecular mechanistic approaches to halt Alzheimer's pathogenesis. *J Cell Physiol* 2019;234:8094-112. [DOI PubMed](#)
16. Piras A, Collin L, Grüninger F, Graff C, Rönnebäck A. Autophagic and lysosomal defects in human tauopathies: analysis of post-mortem brain from patients with familial Alzheimer disease, corticobasal degeneration and progressive supranuclear palsy. *Acta Neuropathol Commun* 2016;4:22. [DOI PubMed PMC](#)
17. Menzies FM, Fleming A, Rubinsztein DC. Compromised autophagy and neurodegenerative diseases. *Nat Rev Neurosci* 2015;16:345-57. [DOI PubMed](#)
18. Nixon RA. The role of autophagy in neurodegenerative disease. *Nat Med* 2013;19:983-97. [DOI PubMed](#)
19. Pickford F, Masliah E, Britschgi M, et al. The autophagy-related protein beclin 1 shows reduced expression in early Alzheimer disease and regulates amyloid beta accumulation in mice. *J Clin Invest* 2008;118:2190-9. [DOI PubMed PMC](#)
20. Shibata M, Lu T, Furuya T, et al. Regulation of intracellular accumulation of mutant Huntingtin by Beclin 1. *J Biol Chem* 2006;281:14474-85. [DOI PubMed](#)
21. Ando K, Brion JP, Stygelbout V, et al. Clathrin adaptor CALM/PICALM is associated with neurofibrillary tangles and is cleaved in Alzheimer's brains. *Acta Neuropathol* 2013;125:861-78. [DOI PubMed](#)
22. Jun G, Naj AC, Beecham GW, et al; Alzheimer's Disease Genetics Consortium. Meta-analysis confirms CR1, CLU, and PICALM as alzheimer disease risk loci and reveals interactions with APOE genotypes. *Arch Neurol* 2010;67:1473-84. [DOI PubMed PMC](#)
23. Moreau K, Fleming A, Imarisio S, et al. PICALM modulates autophagy activity and tau accumulation. *Nat Commun* 2014;5:4998. [DOI PubMed PMC](#)
24. Nixon RA, Yang DS. Autophagy failure in Alzheimer's disease--locating the primary defect. *Neurobiol Dis* 2011;43:38-45. [DOI PubMed PMC](#)
25. Dixit R, Ross JL, Goldman YE, Holzbaur EL. Differential regulation of dynein and kinesin motor proteins by tau. *Science* 2008;319:1086-9. [DOI PubMed PMC](#)
26. Ittner LM, Ke YD, Delerue F, et al. Dendritic function of tau mediates amyloid-beta toxicity in Alzheimer's disease mouse models. *Cell* 2010;142:387-97. [DOI PubMed](#)
27. Pavel M, Imarisio S, Menzies FM, et al. CCT complex restricts neuropathogenic protein aggregation via autophagy. *Nat Commun* 2016;7:13821. [DOI PubMed PMC](#)
28. Brehme M, Voisine C, Rolland T, et al. A chaperome subnetwork safeguards proteostasis in aging and neurodegenerative disease. *Cell Rep* 2014;9:1135-50. [DOI PubMed PMC](#)
29. Lee J, Yu WH, Kumar A, et al. Lysosomal proteolysis and autophagy require presenilin 1 and are disrupted by alzheimer-related PS1 mutations. *Cell* 2010;141:1146-58. [DOI PubMed PMC](#)
30. Lee JK, Jin HK, Park MH, et al. Acid sphingomyelinase modulates the autophagic process by controlling lysosomal biogenesis in Alzheimer's disease. *J Exp Med* 2014;211:1551-70. [DOI PubMed PMC](#)
31. Ejlerskov P, Ashkenazi A, Rubinsztein DC. Genetic enhancement of macroautophagy in vertebrate models of neurodegenerative diseases. *Neurobiol Dis* 2019;122:3-8. [DOI PubMed](#)
32. Jiang T, Yu JT, Zhu XC, et al. Temsirolimus attenuates tauopathy in vitro and in vivo by targeting tau hyperphosphorylation and autophagic clearance. *Neuropharmacology* 2014;85:121-30. [DOI PubMed](#)
33. Ozelik S, Fraser G, Castets P, et al. Rapamycin attenuates the progression of tau pathology in P301S tau transgenic mice. *PLoS One* 2013;8:e62459. [DOI PubMed PMC](#)
34. Spilman P, Podlutska N, Hart MJ, et al. Inhibition of mTOR by rapamycin abolishes cognitive deficits and reduces amyloid-beta levels in a mouse model of Alzheimer's disease. *PLoS One* 2010;5:e9979. [DOI PubMed PMC](#)
35. Berger Z, Ravikumar B, Menzies FM, et al. Rapamycin alleviates toxicity of different aggregate-prone proteins. *Hum Mol Genet* 2006;15:433-42. [DOI PubMed](#)
36. Karabiyik C, Vicinanza M, Rubinsztein DC. AMPK: the energy sensor that regulates autophagy and a potential therapeutic target for neurodegenerative diseases. Non-canonical autophagy. Elsevier; 2021. p. 9-39. [DOI](#)
37. Hansen M, Rubinsztein DC, Walker DW. Publisher correction: autophagy as a promoter of longevity: insights from model organisms. *Nat Rev Mol Cell Biol* 2018;19:611. [DOI PubMed](#)
38. Menzies FM, Fleming A, Caricasole A, et al. Autophagy and neurodegeneration: pathogenic mechanisms and therapeutic opportunities. *Neuron* 2017;93:1015-34. [DOI PubMed](#)
39. Ou Z, Kong X, Sun X, et al. Metformin treatment prevents amyloid plaque deposition and memory impairment in APP/PS1 mice. *Brain Behav Immun* 2018;69:351-63. [DOI PubMed](#)
40. Son SM, Shin HJ, Byun J, et al. Metformin facilitates amyloid- β generation by β - and γ -secretases via autophagy activation. *J Alzheimers Dis* 2016;51:1197-208. [DOI PubMed](#)
41. La Barbera L, Vedele F, Nobili A, et al. Nilotinib restores memory function by preventing dopaminergic neuron degeneration in a mouse model of Alzheimer's Disease. *Prog Neurobiol* 2021;202:102031. [DOI PubMed](#)
42. Koenig AM, Mechanic-Hamilton D, Xie SX, et al. Effects of the insulin sensitizer metformin in Alzheimer disease: pilot data from a

- randomized placebo-controlled crossover study. *Alzheimer Dis Assoc Disord* 2017;31:107-13. DOI PubMed PMC
43. Luchsinger JA, Perez T, Chang H, et al. Metformin in amnesic mild cognitive impairment: results of a pilot randomized placebo controlled clinical trial. *J Alzheimers Dis* 2016;51:501-14. DOI PubMed PMC
 44. DeBosch BJ, Heitmeier MR, Mayer AL, et al. Trehalose inhibits solute carrier 2A (SLC2A) proteins to induce autophagy and prevent hepatic steatosis. *Sci Signal* 2016;9:ra21. DOI PubMed PMC
 45. Renna M, Jimenez-Sanchez M, Sarkar S, Rubinsztein DC. Chemical inducers of autophagy that enhance the clearance of mutant proteins in neurodegenerative diseases. *J Biol Chem* 2010;285:11061-7. DOI PubMed PMC
 46. Schaeffer V, Lavenir I, Ozcelik S, Tolnay M, Winkler DT, Goedert M. Stimulation of autophagy reduces neurodegeneration in a mouse model of human tauopathy. *Brain* 2012;135:2169-77. DOI PubMed PMC
 47. Sarkar S, Floto RA, Berger Z, et al. Lithium induces autophagy by inhibiting inositol monophosphatase. *J Cell Biol* 2005;170:1101-11. DOI PubMed PMC
 48. Cárdenas C, Miller RA, Smith I, et al. Essential regulation of cell bioenergetics by constitutive InsP3 receptor Ca²⁺ transfer to mitochondria. *Cell* 2010;142:270-83. DOI PubMed PMC
 49. Zhang X, Heng X, Li T, et al. Long-term treatment with lithium alleviates memory deficits and reduces amyloid- β production in an aged Alzheimer's disease transgenic mouse model. *J Alzheimers Dis* 2011;24:739-49. DOI PubMed
 50. Forlenza OV, Radanovic M, Talib LL, Gattaz WF. Clinical and biological effects of long-term lithium treatment in older adults with amnesic mild cognitive impairment: randomised clinical trial. *Br J Psychiatry* 2019;215:668-74. DOI PubMed
 51. Williams A, Sarkar S, Cuddon P, et al. Novel targets for Huntington's disease in an mTOR-independent autophagy pathway. *Nat Chem Biol* 2008;4:295-305. DOI PubMed PMC
 52. Lopez A, Lee SE, Wojta K, et al; Tauopathy Genetics Consortium. A152T tau allele causes neurodegeneration that can be ameliorated in a zebrafish model by autophagy induction. *Brain* 2017;140:1128-46. DOI PubMed PMC
 53. Siddiqi FH, Menzies FM, Lopez A, et al. Felodipine induces autophagy in mouse brains with pharmacokinetics amenable to repurposing. *Nat Commun* 2019;10:1817. DOI PubMed PMC
 54. Medeiros R, Kitazawa M, Chabrier MA, et al. Calpain inhibitor A-705253 mitigates Alzheimer's disease-like pathology and cognitive decline in aged 3xTgAD mice. *Am J Pathol* 2012;181:616-25. DOI PubMed
 55. Khandelwal PJ, Herman AM, Hoe HS, Rebeck GW, Moussa CE. Parkin mediates beclin-dependent autophagic clearance of defective mitochondria and ubiquitinated Abeta in AD models. *Hum Mol Genet* 2011;20:2091-102. DOI PubMed PMC
 56. Rocchi A, Yamamoto S, Ting T, et al. A Becn1 mutation mediates hyperactive autophagic sequestration of amyloid oligomers and improved cognition in Alzheimer's disease. *PLoS Genet* 2017;13:e1006962. DOI PubMed PMC
 57. Yang DS, Stavrides P, Saito M, et al. Defective macroautophagic turnover of brain lipids in the TgCRND8 Alzheimer mouse model: prevention by correcting lysosomal proteolytic deficits. *Brain* 2014;137:3300-18. DOI PubMed PMC
 58. Settembre C, Di Malta C, Polito VA, et al. TFEB links autophagy to lysosomal biogenesis. *Science* 2011;332:1429-33. DOI PubMed PMC
 59. Polito VA, Li H, Martini-Stoica H, et al. Selective clearance of aberrant tau proteins and rescue of neurotoxicity by transcription factor EB. *EMBO Mol Med* 2014;6:1142-60. DOI PubMed PMC
 60. Spillantini MG, Schmidt ML, Lee VM, Trojanowski JQ, Jakes R, Goedert M. Alpha-synuclein in Lewy bodies. *Nature* 1997;388:839-40. DOI PubMed
 61. Ross OA, Braithwaite AT, Skipper LM, et al. Genomic investigation of alpha-synuclein multiplication and parkinsonism. *Ann Neurol* 2008;63:743-50. DOI PubMed PMC
 62. Karabiyik C, Lee MJ, Rubinsztein DC. Autophagy impairment in Parkinson's disease. *Essays Biochem* 2017;61:711-20. DOI PubMed
 63. Tanik SA, Schultheiss CE, Volpicelli-Daley LA, Brunden KR, Lee VM. Lewy body-like α -synuclein aggregates resist degradation and impair macroautophagy. *J Biol Chem* 2013;288:15194-210. DOI PubMed PMC
 64. Winslow AR, Chen CW, Corrochano S, et al. α -Synuclein impairs macroautophagy: implications for Parkinson's disease. *J Cell Biol* 2010;190:1023-37. DOI PubMed PMC
 65. Zavodszky E, Seaman MN, Moreau K, et al. Mutation in VPS35 associated with Parkinson's disease impairs WASH complex association and inhibits autophagy. *Nat Commun* 2014;5:3828. DOI PubMed PMC
 66. Sarkar S, Olsen AL, Sygnecka K, Lohr KM, Feany MB. α -synuclein impairs autophagosome maturation through abnormal actin stabilization. *PLoS Genet* 2021;17:e1009359. DOI PubMed PMC
 67. Qiao L, Hamamichi S, Caldwell KA, et al. Lysosomal enzyme cathepsin D protects against alpha-synuclein aggregation and toxicity. *Mol Brain* 2008;1:17. DOI PubMed PMC
 68. Dodson MW, Zhang T, Jiang C, Chen S, Guo M. Roles of the Drosophila LRRK2 homolog in Rab7-dependent lysosomal positioning. *Hum Mol Genet* 2012;21:1350-63. DOI PubMed PMC
 69. Schöndorf DC, Aureli M, McAllister FE, et al. iPSC-derived neurons from GBA1-associated Parkinson's disease patients show autophagic defects and impaired calcium homeostasis. *Nat Commun* 2014;5:4028. DOI PubMed
 70. Bento CF, Ashkenazi A, Jimenez-Sanchez M, Rubinsztein DC. The Parkinson's disease-associated genes ATP13A2 and SYT11 regulate autophagy via a common pathway. *Nat Commun* 2016;7:11803. DOI PubMed PMC
 71. Borsche M, König IR, Delcambre S, et al. Mitochondrial damage-associated inflammation highlights biomarkers in PRKN/PINK1 parkinsonism. *Brain* 2020;143:3041-51. DOI PubMed PMC
 72. Webb JL, Ravikumar B, Atkins J, Skepper JN, Rubinsztein DC. Alpha-Synuclein is degraded by both autophagy and the proteasome. *J Biol Chem* 2003;278:25009-13. DOI PubMed

73. Dehay B, Bové J, Rodríguez-Muela N, et al. Pathogenic lysosomal depletion in Parkinson's disease. *J Neurosci* 2010;30:12535-44. DOI PubMed PMC
74. Malagelada C, Jin ZH, Jackson-Lewis V, Przedborski S, Greene LA. Rapamycin protects against neuron death in in vitro and in vivo models of Parkinson's disease. *J Neurosci* 2010;30:1166-75. DOI PubMed PMC
75. Crews L, Spencer B, Desplats P, et al. Selective molecular alterations in the autophagy pathway in patients with Lewy body disease and in models of alpha-synucleinopathy. *PLoS One* 2010;5:e9313. DOI PubMed PMC
76. Sarkar S, Perlstein EO, Imarisio S, et al. Small molecules enhance autophagy and reduce toxicity in Huntington's disease models. *Nat Chem Biol* 2007;3:331-8. DOI PubMed PMC
77. Dulovic M, Jovanovic M, Xilouri M, et al. The protective role of AMP-activated protein kinase in alpha-synuclein neurotoxicity in vitro. *Neurobiol Dis* 2014;63:1-11. DOI PubMed
78. Ng CH, Guan MS, Koh C, et al. AMP kinase activation mitigates dopaminergic dysfunction and mitochondrial abnormalities in Drosophila models of Parkinson's disease. *J Neurosci* 2012;32:14311-7. DOI PubMed PMC
79. Hebron ML, Lonskaya I, Moussa CE. Nilotinib reverses loss of dopamine neurons and improves motor behavior via autophagic degradation of α -synuclein in Parkinson's disease models. *Hum Mol Genet* 2013;22:3315-28. DOI PubMed PMC
80. Simuni T, Fiske B, Merchant K, et al. NILO-PD: a phase 2A study of nilotinib in patients with advanced and early parkinson's disease: study design and status update (P3.8-035). *Neurology* 2019:92. DOI
81. Ferretta A, Gaballo A, Tanzarella P, et al. Effect of resveratrol on mitochondrial function: implications in parkin-associated familial Parkinson's disease. *Biochim Biophys Acta* 2014;1842:902-15. DOI PubMed
82. Park SJ, Ahmad F, Philp A, et al. Resveratrol ameliorates aging-related metabolic phenotypes by inhibiting cAMP phosphodiesterases. *Cell* 2012;148:421-33. DOI PubMed PMC
83. Park D, Jeong H, Lee MN, et al. Resveratrol induces autophagy by directly inhibiting mTOR through ATP competition. *Sci Rep* 2016;6:21772. DOI PubMed PMC
84. Lieu CA, Dewey CM, Chinta SJ, et al. Lithium prevents parkinsonian behavioral and striatal phenotypes in an aged parkin mutant transgenic mouse model. *Brain Res* 2014;1591:111-7. DOI PubMed PMC
85. Xiong N, Jia M, Chen C, et al. Potential autophagy enhancers attenuate rotenone-induced toxicity in SH-SY5Y. *Neuroscience* 2011;199:292-302. DOI PubMed
86. Sarkar S, Davies JE, Huang Z, Tunnacliffe A, Rubinsztein DC. Trehalose, a novel mTOR-independent autophagy enhancer, accelerates the clearance of mutant huntingtin and alpha-synuclein. *J Biol Chem* 2007;282:5641-52. DOI PubMed
87. Tanji K, Miki Y, Maruyama A, et al. Trehalose intake induces chaperone molecules along with autophagy in a mouse model of Lewy body disease. *Biochem Biophys Res Commun* 2015;465:746-52. DOI PubMed
88. Jimenez-Sanchez M, Thomson F, Zavodszky E, Rubinsztein DC. Autophagy and polyglutamine diseases. *Prog Neurobiol* 2012;97:67-82. DOI PubMed PMC
89. McLoughlin HS, Moore LR, Paulson HL. Pathogenesis of SCA3 and implications for other polyglutamine diseases. *Neurobiol Dis* 2020;134:104635. DOI PubMed PMC
90. Kratter IH, Finkbeiner S. PolyQ disease: too many Qs, too much function? *Neuron* 2010;67:897-9. DOI PubMed PMC
91. Cortes CJ, La Spada AR. Autophagy in polyglutamine disease: imposing order on disorder or contributing to the chaos? *Mol Cell Neurosci* 2015;66:53-61. DOI PubMed PMC
92. Macdonald M. A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington's disease chromosomes. *Cell* 1993;72:971-83. DOI PubMed
93. Ravikumar B, Berger Z, Vacher C, O'Kane CJ, Rubinsztein DC. Rapamycin pre-treatment protects against apoptosis. *Hum Mol Genet* 2006;15:1209-16. DOI PubMed
94. Sapp E, Schwarz C, Chase K, et al. Huntingtin localization in brains of normal and Huntington's disease patients. *Ann Neurol* 1997;42:604-12. DOI PubMed
95. Rubinsztein DC. Lessons from animal models of Huntington's disease. *Trends in Genetics* 2002;18:202-9. DOI PubMed
96. Ashkenazi A, Bento CF, Ricketts T, et al. Polyglutamine tracts regulate beclin 1-dependent autophagy. *Nature* 2017;545:108-11. DOI PubMed PMC
97. Martinez-Vicente M, Tallozy Z, Wong E, et al. Cargo recognition failure is responsible for inefficient autophagy in Huntington's disease. *Nat Neurosci* 2010;13:567-76. DOI PubMed PMC
98. Ochaba J, Lukacsovich T, Csikos G, et al. Potential function for the Huntingtin protein as a scaffold for selective autophagy. *Proc Natl Acad Sci U S A* 2014;111:16889-94. DOI PubMed PMC
99. Rui YN, Xu Z, Patel B, et al. Huntingtin functions as a scaffold for selective macroautophagy. *Nat Cell Biol* 2015;17:262-75. DOI PubMed PMC
100. Wong YC, Holzbaur EL. The regulation of autophagosome dynamics by huntingtin and HAP1 is disrupted by expression of mutant huntingtin, leading to defective cargo degradation. *J Neurosci* 2014;34:1293-305. DOI PubMed PMC
101. Ravikumar B, Acevedo-Arozena A, Imarisio S, et al. Dynein mutations impair autophagic clearance of aggregate-prone proteins. *Nat Genet* 2005;37:771-6. DOI PubMed
102. Hinckelmann MV, Zala D, Saudou F. Releasing the brake: restoring fast axonal transport in neurodegenerative disorders. *Trends Cell Biol* 2013;23:634-43. DOI PubMed
103. del Toro D, Alberch J, Lázaro-Díéguez F, et al. Mutant huntingtin impairs post-Golgi trafficking to lysosomes by delocalizing optineurin/Rab8 complex from the Golgi apparatus. *Mol Biol Cell* 2009;20:1478-92. DOI PubMed PMC
104. Subramaniam S, Sixt KM, Barrow R, Snyder SH. Rhes, a striatal specific protein, mediates mutant-huntingtin cytotoxicity. *Science*

- 2009;324:1327-30. DOI PubMed PMC
105. Mealer RG, Murray AJ, Shahani N, Subramaniam S, Snyder SH. Rhes, a striatal-selective protein implicated in Huntington disease, binds beclin-1 and activates autophagy. *J Biol Chem* 2014;289:3547-54. DOI PubMed PMC
 106. Metzger S, Saukko M, Van Che H, et al. Age at onset in Huntington's disease is modified by the autophagy pathway: implication of the V471A polymorphism in Atg7. *Hum Genet* 2010;128:453-9. DOI PubMed
 107. Franco-Iborra S, Plaza-Zabala A, Montpeyo M, Sebastian D, Vila M, Martínez-Vicente M. Mutant HTT (huntingtin) impairs mitophagy in a cellular model of Huntington disease. *Autophagy* 2021;17:672-89. DOI PubMed PMC
 108. Koga H, Martínez-Vicente M, Arias E, Kaushik S, Sulzer D, Cuervo AM. Constitutive upregulation of chaperone-mediated autophagy in Huntington's disease. *J Neurosci* 2011;31:18492-505. DOI PubMed PMC
 109. Ravikumar B, Berger Z, Vacher C, O'Kane CJ, Rubinsztein DC. Rapamycin pre-treatment protects against apoptosis. *Hum Mol Genet* 2006;15:1209-16. DOI PubMed
 110. Zheng S, Clabough EB, Sarkar S, Futter M, Rubinsztein DC, Zeitlin SO. Deletion of the huntingtin polyglutamine stretch enhances neuronal autophagy and longevity in mice. *PLoS Genet* 2010;6:e1000838. DOI PubMed PMC
 111. Menzies FM, Hourez R, Imarisio S, et al. Puromycin-sensitive aminopeptidase protects against aggregation-prone proteins via autophagy. *Hum Mol Genet* 2010;19:4573-86. DOI PubMed PMC
 112. Sarkar S, Krishna G, Imarisio S, Saiki S, O'Kane CJ, Rubinsztein DC. A rational mechanism for combination treatment of Huntington's disease using lithium and rapamycin. *Hum Mol Genet* 2008;17:170-8. DOI PubMed
 113. Jiang W, Wei W, Gaertig MA, Li S, Li XJ. Therapeutic effect of berberine on Huntington's disease transgenic mouse model. *PLoS One* 2015;10:e0134142. DOI PubMed PMC
 114. Rose C, Menzies FM, Renna M, et al. Rilmenidine attenuates toxicity of polyglutamine expansions in a mouse model of Huntington's disease. *Hum Mol Genet* 2010;19:2144-53. DOI PubMed PMC
 115. Sanchis A, Garcia-Gimeno MA, Cañada-Martínez AJ, et al. Metformin treatment reduces motor and neuropsychiatric phenotypes in the zQ175 mouse model of Huntington disease. *Exp Mol Med* 2019;51:1-16. DOI PubMed PMC
 116. Menzies FM, Garcia-Arencibia M, Imarisio S, et al. Calpain inhibition mediates autophagy-dependent protection against polyglutamine toxicity. *Cell Death Differ* 2015;22:433-44. DOI PubMed PMC
 117. Fox LM, Kim K, Johnson CW, et al. Huntington's disease pathogenesis is modified in vivo by ALFY/WDFY3 and selective macroautophagy. *Neuron* 2020;105:813-821.e6. DOI PubMed PMC
 118. Filimonenko M, Isakson P, Finley KD, et al. The selective macroautophagic degradation of aggregated proteins requires the PI3P-binding protein Alf1. *Mol Cell* 2010;38:265-79. DOI PubMed PMC
 119. Vodicka P, Chase K, Iuliano M, et al. Autophagy Activation by transcription factor EB (TFEB) in striatum of HDQ175/Q7 mice. *J Huntingtons Dis* 2016;5:249-60. DOI PubMed PMC
 120. Qi L, Zhang XD, Wu JC, et al. The role of chaperone-mediated autophagy in huntingtin degradation. *PLoS One* 2012;7:e46834. DOI PubMed PMC
 121. Jeong H, Then F, Melia TJ Jr, et al. Acetylation targets mutant huntingtin to autophagosomes for degradation. *Cell* 2009;137:60-72. DOI PubMed PMC
 122. Son SM, Park SJ, Fernandez-Estevez M, Rubinsztein DC. Autophagy regulation by acetylation-implications for neurodegenerative diseases. *Exp Mol Med* 2021;53:30-41. DOI PubMed PMC
 123. van Es MA, Hardiman O, Chio A, et al. Amyotrophic lateral sclerosis. *Lancet* 2017;390:2084-98. DOI PubMed
 124. Robberecht W, Philips T. The changing scene of amyotrophic lateral sclerosis. *Nat Rev Neurosci* 2013;14:248-64. DOI PubMed
 125. van Zundert B, Peuscher MH, Hynynen M, et al. Neonatal neuronal circuitry shows hyperexcitable disturbance in a mouse model of the adult-onset neurodegenerative disease amyotrophic lateral sclerosis. *J Neurosci* 2008;28:10864-74. DOI PubMed PMC
 126. Amendola J, Verrier B, Roubertoux P, Durand J. Altered sensorimotor development in a transgenic mouse model of amyotrophic lateral sclerosis. *Eur J Neurosci* 2004;20:2822-6. DOI PubMed
 127. Williamson TL, Cleveland DW. Slowing of axonal transport is a very early event in the toxicity of ALS-linked SOD1 mutants to motor neurons. *Nat Neurosci* 1999;2:50-6. DOI PubMed
 128. Pandya VA, Patani R. Decoding the relationship between ageing and amyotrophic lateral sclerosis: a cellular perspective. *Brain* 2020;143:1057-72. DOI PubMed PMC
 129. Jacob JM. Lumbar motor neuron size and number is affected by age in male F344 rats. *Mech Ageing Dev* 1998;106:205-16. DOI PubMed
 130. Tomlinson B, Irving D. The numbers of limb motor neurons in the human lumbosacral cord throughout life. *J Neurol Sci* 1977;34:213-9. DOI PubMed
 131. Moldovan M, Rosberg MR, Alvarez S, Klein D, Martini R, Krarup C. Aging-associated changes in motor axon voltage-gated Na(+) channel function in mice. *Neurobiol Aging* 2016;39:128-39. DOI PubMed
 132. Rygiel KA, Grady JP, Turnbull DM. Respiratory chain deficiency in aged spinal motor neurons. *Neurobiol Aging* 2014;35:2230-8. DOI PubMed PMC
 133. Fernando R, Castro JP, Flore T, Deubel S, Grune T, Ott C. Age-related maintenance of the autophagy-lysosomal system is dependent on skeletal muscle type. *Oxid Med Cell Longev* 2020;2020:4908162. DOI PubMed PMC
 134. Carnio S, LoVerso F, Baraibar MA, et al. Autophagy impairment in muscle induces neuromuscular junction degeneration and precocious aging. *Cell Rep* 2014;8:1509-21. DOI PubMed PMC
 135. Amin A, Perera ND, Beart PM, Turner BJ, Shabanpoor F. Amyotrophic lateral sclerosis and autophagy: dysfunction and therapeutic targeting. *Cells* 2020;9:2413. DOI PubMed PMC

136. Sundaramoorthy V, Walker AK, Tan V, et al. Defects in optineurin- and myosin VI-mediated cellular trafficking in amyotrophic lateral sclerosis. *Hum Mol Genet* 2015;24:3830-46. DOI PubMed
137. Wild P, Farhan H, McEwan DG, et al. Phosphorylation of the autophagy receptor optineurin restricts Salmonella growth. *Science* 2011;333:228-33. DOI PubMed PMC
138. Freischmidt A, Wieland T, Richter B, et al. Haploinsufficiency of TBK1 causes familial ALS and fronto-temporal dementia. *Nat Neurosci* 2015;18:631-6. DOI PubMed
139. de Majo M, Topp SD, Smith BN, et al. ALS-associated missense and nonsense TBK1 mutations can both cause loss of kinase function. *Neurobiol Aging* 2018;71:266.e1-266.e10. DOI PubMed PMC
140. Fecto F, Yan J, Vemula SP, et al. SQSTM1 mutations in familial and sporadic amyotrophic lateral sclerosis. *Arch Neurol* 2011;68:1440-6. DOI PubMed
141. Ber I, Camuzat A, Guerreiro R, et al; French Clinical and Genetic Research Network on FTD/FTD-ALS. SQSTM1 mutations in French patients with frontotemporal dementia or frontotemporal dementia with amyotrophic lateral sclerosis. *JAMA Neurol* 2013;70:1403-10. DOI PubMed PMC
142. Rubino E, Rainero I, Chiò A, et al; TODEM Study Group. SQSTM1 mutations in frontotemporal lobar degeneration and amyotrophic lateral sclerosis. *Neurology* 2012;79:1556-62. DOI PubMed PMC
143. Goode A, Butler K, Long J, et al. Defective recognition of LC3B by mutant SQSTM1/p62 implicates impairment of autophagy as a pathogenic mechanism in ALS-FTLD. *Autophagy* 2016;12:1094-104. DOI PubMed PMC
144. Momeni P, Bell J, Duckworth J, et al. Sequence analysis of all identified open reading frames on the frontal temporal dementia haplotype on chromosome 3 fails to identify unique coding variants except in CHMP2B. *Neurosci Lett* 2006;410:77-9. DOI PubMed
145. Lee JA, Beigneux A, Ahmad ST, Young SG, Gao FB. ESCRT-III dysfunction causes autophagosome accumulation and neurodegeneration. *Curr Biol* 2007;17:1561-7. DOI PubMed
146. Papadopoulos C, Kirchner P, Bug M, et al. VCP/p97 cooperates with YOD1, UBXD1 and PLAA to drive clearance of ruptured lysosomes by autophagy. *EMBO J* 2017;36:135-50. DOI PubMed PMC
147. Buchan JR, Kolaitis RM, Taylor JP, Parker R. Eukaryotic stress granules are cleared by autophagy and Cdc48/VCP function. *Cell* 2013;153:1461-74. DOI PubMed PMC
148. Hill SM, Wrobel L, Ashkenazi A, et al. VCP/p97 regulates Beclin-1-dependent autophagy initiation. *Nat Chem Biol* 2021;17:448-55. DOI PubMed
149. Casterton RL, Hunt RJ, Fanto M. Pathomechanism heterogeneity in the amyotrophic lateral sclerosis and frontotemporal dementia disease spectrum: providing focus through the lens of autophagy. *J Mol Biol* 2020;432:2692-713. DOI PubMed
150. Zou ZY, Zhou ZR, Che CH, Liu CY, He RL, Huang HP. Genetic epidemiology of amyotrophic lateral sclerosis: a systematic review and meta-analysis. *J Neurol Neurosurg Psychiatry* 2017;88:540-9. DOI PubMed
151. DeJesus-Hernandez M, Mackenzie IR, Boeve BF, et al. Expanded GGGGCC hexanucleotide repeat in noncoding region of C9ORF72 causes chromosome 9p-linked FTD and ALS. *Neuron* 2011;72:245-56. DOI PubMed PMC
152. Renton AE, Majounie E, Waite A, et al; ITALSGEN Consortium. A hexanucleotide repeat expansion in C9ORF72 is the cause of chromosome 9p21-linked ALS-FTD. *Neuron* 2011;72:257-68. DOI PubMed PMC
153. Mori K, Weng SM, Arzberger T, et al. The C9orf72 GGGGCC repeat is translated into aggregating dipeptide-repeat proteins in FTLD/ALS. *Science* 2013;339:1335-8. DOI PubMed
154. Cristofani R, Crippa V, Vezzoli G, et al. The small heat shock protein B8 (HSPB8) efficiently removes aggregating species of dipeptides produced in C9ORF72-related neurodegenerative diseases. *Cell Stress Chaperones* 2018;23:1-12. DOI PubMed PMC
155. Balendra R, Isaacs AM. C9orf72-mediated ALS and FTD: multiple pathways to disease. *Nat Rev Neurol* 2018;14:544-58. DOI PubMed PMC
156. Boivin M, Pfister V, Gaucherot A, et al. Reduced autophagy upon C9ORF72 loss synergizes with dipeptide repeat protein toxicity in G4C2 repeat expansion disorders. *EMBO J* 2020;39:e100574. DOI PubMed PMC
157. Lee EB, Lee VM, Trojanowski JQ. Gains or losses: molecular mechanisms of TDP43-mediated neurodegeneration. *Nat Rev Neurosci* 2011;13:38-50. DOI PubMed PMC
158. Cascella R, Fani G, Capitini C, et al. Quantitative assessment of the degradation of aggregated TDP-43 mediated by the ubiquitin proteasome system and macroautophagy. *FASEB J* 2017;31:5609-24. DOI PubMed
159. Scotter EL, Vance C, Nishimura AL, et al. Differential roles of the ubiquitin proteasome system and autophagy in the clearance of soluble and aggregated TDP-43 species. *J Cell Sci* 2014;127:1263-78. DOI PubMed PMC
160. Bose JK, Huang CC, Shen CK. Regulation of autophagy by neuropathological protein TDP-43. *J Biol Chem* 2011;286:44441-8. DOI PubMed PMC
161. Kabuta T, Suzuki Y, Wada K. Degradation of amyotrophic lateral sclerosis-linked mutant Cu,Zn-superoxide dismutase proteins by macroautophagy and the proteasome. *J Biol Chem* 2006;281:30524-33. DOI PubMed
162. Deng HX, Zhai H, Bigio EH, et al. FUS-immunoreactive inclusions are a common feature in sporadic and non-SOD1 familial amyotrophic lateral sclerosis. *Ann Neurol* 2010;67:739-48. DOI PubMed PMC
163. Madruga E, Maestro I, Martínez A. Mitophagy modulation, a new player in the race against ALS. *Int J Mol Sci* 2021;22:740. DOI PubMed PMC
164. Brown DG, Shorter J, Wobst HJ. Emerging small-molecule therapeutic approaches for amyotrophic lateral sclerosis and frontotemporal dementia. *Bioorg Med Chem Lett* 2020;30:126942. DOI PubMed
165. Mandrioli J, D'Amico R, Zucchi E, et al; RAP-ALS investigators group. Rapamycin treatment for amyotrophic lateral sclerosis: Protocol for a phase II randomized, double-blind, placebo-controlled, multicenter, clinical trial (RAP-ALS trial). *Medicine*

- (Baltimore) 2018;97:e11119. DOI PubMed PMC
166. Wang IF, Guo BS, Liu YC, et al. Autophagy activators rescue and alleviate pathogenesis of a mouse model with proteinopathies of the TAR DNA-binding protein 43. *Proc Natl Acad Sci U S A* 2012;109:15024-9. DOI PubMed PMC
 167. Cheng CW, Lin MJ, Shen CK. Rapamycin alleviates pathogenesis of a new Drosophila model of ALS-TDP. *J Neurogenet* 2015;29:59-68. DOI PubMed
 168. Zhang X, Li L, Chen S, et al. Rapamycin treatment augments motor neuron degeneration in SOD1(G93A) mouse model of amyotrophic lateral sclerosis. *Autophagy* 2011;7:412-25. DOI PubMed
 169. Li Y, Guo Y, Wang X, et al. Trehalose decreases mutant SOD1 expression and alleviates motor deficiency in early but not end-stage amyotrophic lateral sclerosis in a SOD1-G93A mouse model. *Neuroscience* 2015;298:12-25. DOI PubMed
 170. Fornai F, Longone P, Cafaro L, et al. Lithium delays progression of amyotrophic lateral sclerosis. *Proc Natl Acad Sci U S A* 2008;105:16404-7. DOI PubMed PMC
 171. in patients with amyotrophic lateral sclerosis (LiCALS): a phase 3 multicentre, randomised, double-blind, placebo-controlled trial. *Lancet Neurol* 2013;12:339-45. DOI PubMed PMC
 172. Chiò A, Borghero G, Calvo A, et al; LITALS Study Group. Lithium carbonate in amyotrophic lateral sclerosis: lack of efficacy in a dose-finding trial. *Neurology* 2010;75:619-25. DOI PubMed
 173. Aggarwal SP, Zinman L, Simpson E, et al. Safety and efficacy of lithium in combination with riluzole for treatment of amyotrophic lateral sclerosis: a randomised, double-blind, placebo-controlled trial. *Lancet Neurol* 2010;9:481-8. DOI PubMed PMC
 174. Verstraete E, Veldink JH, Huisman MH, et al. Lithium lacks effect on survival in amyotrophic lateral sclerosis: a phase IIb randomised sequential trial. *J Neurol Neurosurg Psychiatry* 2012;83:557-64. DOI PubMed
 175. Eloheid A, Libard S, Leino M, Popova SN, Alafuzoff I. Altered proteins in the aging brain. *J Neuropathol Exp Neurol* 2016;75:316-25. DOI PubMed PMC
 176. Hansen M, Chandra A, Mitic LL, Onken B, Driscoll M, Kenyon C. A role for autophagy in the extension of lifespan by dietary restriction in *C. elegans*. *PLoS Genet* 2008;4:e24. DOI PubMed PMC
 177. Demontis F, Perrimon N. FOXO/4E-BP signaling in Drosophila muscles regulates organism-wide proteostasis during aging. *Cell* 2010;143:813-25. DOI PubMed PMC
 178. Sarkis G, Ashcom J, Hawdon J, Jacobson L. Decline in protease activities with age in the nematode *Caenorhabditis elegans*. *Mech Ageing Dev* 1988;45:191-201. DOI PubMed
 179. Simonsen A, Cumming RC, Brech A, Isakson P, Schubert DR, Finley KD. Promoting basal levels of autophagy in the nervous system enhances longevity and oxidant resistance in adult Drosophila. *Autophagy* 2008;4:176-84. DOI PubMed
 180. Lipinski MM, Zheng B, Lu T, et al. Genome-wide analysis reveals mechanisms modulating autophagy in normal brain aging and in Alzheimer's disease. *Proc Natl Acad Sci U S A* 2010;107:14164-9. DOI PubMed PMC
 181. Kaushik S, Arias E, Kwon H, et al. Loss of autophagy in hypothalamic POMC neurons impairs lipolysis. *EMBO Rep* 2012;13:258-65. DOI PubMed PMC
 182. Cuervo AM, Dice JF. Age-related decline in chaperone-mediated autophagy. *J Biol Chem* 2000;275:31505-13. DOI PubMed
 183. Chang JT, Kumsta C, Hellman AB, Adams LM, Hansen M. Spatiotemporal regulation of autophagy during *Caenorhabditis elegans* aging. *Elife* 2017;6:1-23. DOI PubMed PMC
 184. Wilhelm T, Byrne J, Medina R, et al. Neuronal inhibition of the autophagy nucleation complex extends lifespan in post-reproductive *C. elegans*. *Genes Dev* 2017;31:1561-72. DOI PubMed PMC
 185. Roso A. Ageing-related changes in the in vivo function of rat liver macroautophagy and proteolysis. *Exp Gerontol* 2003;38:519-27. DOI PubMed
 186. Donati A, Cavallini G, Paradiso C, et al. Age-related changes in the regulation of autophagic proteolysis in rat isolated hepatocytes. *J Gerontol A Biol Sci Med Sci* 2001;56:B288-93. DOI PubMed
 187. Terman A. The effect of age on formation and elimination of autophagic vacuoles in mouse hepatocytes. *Gerontology* 1995;41 Suppl 2:319-26. DOI PubMed
 188. Meléndez A, Tallóczy Z, Seaman M, Eskelinen EL, Hall DH, Levine B. Autophagy genes are essential for dauer development and life-span extension in *C. elegans*. *Science* 2003;301:1387-91. DOI PubMed
 189. Juhász G, Erdi B, Sass M, Neufeld TP. Atg7-dependent autophagy promotes neuronal health, stress tolerance, and longevity but is dispensable for metamorphosis in Drosophila. *Genes Dev* 2007;21:3061-6. DOI PubMed PMC
 190. Ma Q, Qiang J, Gu P, Wang Y, Geng Y, Wang M. Age-related autophagy alterations in the brain of senescence accelerated mouse prone 8 (SAMP8) mice. *Exp Gerontol* 2011;46:533-41. DOI PubMed
 191. Cavallini G, Donati A, Gori Z, Pollera M, Bergamini E. The protection of rat liver autophagic proteolysis from the age-related decline co-varies with the duration of anti-ageing food restriction. *Experimental Gerontology* 2001;36:497-506. DOI PubMed
 192. Donati A, Ventrucci A, Cavallini G, et al. In vivo effect of an antilipolytic drug (3,5'-dimethylpyrazole) on autophagic proteolysis and autophagy-related gene expression in rat liver. *Biochem Biophys Res Commun* 2008;366:786-92. DOI PubMed
 193. Bai H, Kang P, Hernandez AM, Tatar M. Activin signaling targeted by insulin/dFOXO regulates aging and muscle proteostasis in Drosophila. *PLoS Genet* 2013;9:e1003941. DOI PubMed PMC
 194. Pyo JO, Yoo SM, Ahn HH, et al. Overexpression of Atg5 in mice activates autophagy and extends lifespan. *Nat Commun* 2013;4:2300. DOI PubMed PMC
 195. Lapierre LR, De Magalhaes Filho CD, McQuary PR, et al. The TFEB orthologue HLH-30 regulates autophagy and modulates longevity in *Caenorhabditis elegans*. *Nat Commun* 2013;4:2267. DOI PubMed PMC
 196. Fernández ÁF, Sebti S, Wei Y, et al. Disruption of the beclin 1-BCL2 autophagy regulatory complex promotes longevity in mice.

- Nature* 2018;558:136-40. DOI PubMed PMC
197. Sun N, Youle RJ, Finkel T. The mitochondrial basis of aging. *Mol Cell* 2016;61:654-66. DOI PubMed PMC
 198. Green DR, Galluzzi L, Kroemer G. Mitochondria and the autophagy-inflammation-cell death axis in organismal aging. *Science* 2011;333:1109-12. DOI PubMed PMC
 199. Rana A, Oliveira MP, Khamoui AV, et al. Promoting Drp1-mediated mitochondrial fission in midlife prolongs healthy lifespan of *Drosophila melanogaster*. *Nat Commun* 2017;8:448. DOI PubMed PMC
 200. Schiavi A, Maglioni S, Palikaras K, et al. Iron-starvation-induced mitophagy mediates lifespan extension upon mitochondrial stress in *C. elegans*. *Curr Biol* 2015;25:1810-22. DOI PubMed
 201. Ryu D, Mouchiroud L, Andreux PA, et al. Urolithin A induces mitophagy and prolongs lifespan in *C. elegans* and increases muscle function in rodents. *Nat Med* 2016;22:879-88. DOI PubMed
 202. Matecic M, Smith DL, Pan X, et al. A microarray-based genetic screen for yeast chronological aging factors. *PLoS Genet* 2010;6:e1000921. DOI PubMed PMC
 203. Alvers AL, Fishwick LK, Wood MS, et al. Autophagy and amino acid homeostasis are required for chronological longevity in *Saccharomyces cerevisiae*. *Aging Cell* 2009;8:353-69. DOI PubMed PMC
 204. Jia K, Levine B. Autophagy is required for dietary restriction-mediated life span extension in *C. elegans*. *Autophagy* 2007;3:597-9. DOI PubMed
 205. Hars ES, Qi H, Ryazanov AG, et al. Autophagy regulates ageing in *C. elegans*. *Autophagy* 2007;3:93-5. DOI PubMed
 206. Harrison DE, Strong R, Sharp ZD, et al. Rapamycin fed late in life extends lifespan in genetically heterogeneous mice. *Nature* 2009;460:392-5. DOI PubMed PMC
 207. Bjedov I, Toivonen JM, Kerr F, et al. Mechanisms of life span extension by rapamycin in the fruit fly *Drosophila melanogaster*. *Cell Metab* 2010;11:35-46. DOI PubMed PMC
 208. Laplante M, Sabatini DM. mTOR signaling in growth control and disease. *Cell* 2012;149:274-93. DOI PubMed PMC